

PLEASE READ:


Important reminders about injecting TRYNGOLZA



TRYNGOLZA is for adults with very high triglycerides, also known as severe hypertriglyceridemia, and adults with familial chylomicronemia syndrome or FCS, used along with diet. The tips below can help you feel confident when administering your monthly injection. These are not all the steps for injecting TRYNGOLZA. For additional information, please see complete [Instructions for Use](#) for TRYNGOLZA and/or talk to your doctor.


Three easy steps for injecting TRYNGOLZA

Place




Remove the clear cap by pulling it straight off. Place the orange needle shield at a 90-degree angle against the skin. **Make sure you can see the viewing window.**

Push



Push firmly and hold the autoinjector straight against the skin. You will hear a click as the injection starts. **You may hear a second click. This is normal. The procedure is not finished.**

Hold



Continue to **hold the autoinjector firmly against the skin for 10 seconds** to ensure the full dose has been given. Remove the autoinjector by lifting straight up.



TRYNGOLZA is available in 80 mg and 50 mg doses. The label on your autoinjector may look different, depending on the dose your healthcare provider prescribes.



- Do not push the orange needle shield against the hand or finger
- Do not move, turn, or change the angle of the autoinjector during the injection
- Check that the orange plunger rod filled the entire viewing window to ensure the full dose was delivered
- Put the used autoinjector in a sharps container after use

[Click to watch an instructional video](#)

SELECT IMPORTANT SAFETY INFORMATION & INDICATIONS

Do not use TRYNGOLZA if you have had a serious allergic reaction to TRYNGOLZA or any of the ingredients in TRYNGOLZA.

Please see Important Safety Information throughout, and full [Prescribing Information](#) and [Patient Information](#) for TRYNGOLZA.



[Click to watch an instructional video](#) 

For more details on injecting TRYNGOLZA, contact Ionis Every Step[™] support at 1-844-789-8744.

SELECT IMPORTANT SAFETY INFORMATION & INDICATIONS

Before using TRYNGOLZA, tell your healthcare provider about all of your medical conditions, including if you: are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. It is not known if TRYNGOLZA can harm your unborn baby, or if it passes into your breast milk and if it can harm your baby.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of TRYNGOLZA?

TRYNGOLZA can cause side effects including allergic reactions that may be serious. Allergic reactions can include redness of the skin, red itchy bumps (hives), swelling of the face, chills or trouble breathing. Stop taking TRYNGOLZA and call your healthcare provider or get emergency help right away if you have any of these symptoms.

TRYNGOLZA can cause increases in liver enzymes and fat stored inside the liver. Your healthcare provider may do liver tests before you start taking TRYNGOLZA or if there is an increase in your dose. Tell your healthcare provider right away if you have the following symptoms: feeling tired or weak, right upper stomach discomfort, yellowing of the skin and eyes, loss of appetite, or dark colored urine.

The most common side effects of TRYNGOLZA in people with FCS include injection site reactions (such as redness, itching, rash, or pain at the injection site), decreased platelet count (blood cells that help to clot blood), and joint pain or stiffness.

The most common side effects of TRYNGOLZA in people with sHTG include injection site reactions (such as redness, itching, rash, or pain at the injection site) and increased liver enzymes.

These are not all the possible side effects of TRYNGOLZA. Tell your healthcare provider or treatment team about any side effect you may have.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/safety/medwatch, or call 1-800-FDA-1088.

INDICATIONS

TRYNGOLZA (olezarsen) is a prescription medicine used along with diet to:

- Reduce triglycerides (fat in the blood) in the treatment of adults with a condition that keeps the body from breaking down fats called familial chylomicronemia syndrome (FCS).
- Reduce triglycerides and reduce the risk of acute inflammation of your pancreas (pancreatitis) in the treatment of adults with a condition marked by very high levels of triglycerides in the blood called severe hypertriglyceridemia (sHTG).

It is not known if TRYNGOLZA is safe and effective in children.

Please see Important Safety Information throughout, and full Prescribing Information and Patient Information for TRYNGOLZA.